arXiv:math/0602534v2 [math.AG] 15 Oct 2007

EULER CHARACTERISTIC OF PRIMITIVE T-HYPERSURFACES AND MAXIMAL SURFACES

BENOIT BERTRAND

Abstract. Viro method plays an important role in the study of topology of real algebraic hypersurfaces. The T-primitive hypersurfaces we study here appear as the result of Viro's combinatorial patchworking when one starts with a primitive triangulation. We show that the Euler characteristic of the real part of such a hypersurface of even dimension is equal to the signature of its complex part. We use this result to prove the existence of maximal surfaces in some three-dimensional toric varieties, namely those corresponding to Nakajima polytopes. In fact, these results belong to the field of tropical geometry and we explain how they can be understood tropically.

Contents

Introduction 2
1. Preliminaries 3
1.1. Toric varieties 3
1.2. Combinatorial patchworking 3
1.3. Tropical hypersurfaces 5
1.4. Tropical Statements 9
1.5. Some facts about triangulations of lattice polytopes 10
1.6. Danilov and Khovanskii formulae 11
2. Primitive T-hypersurfaces 12
2.1. Statement 12
2.2. Proof of Theorem 2.1 13
3. Nakajima polytopes and M-surfaces 17
3.1. Newton polytopes without maximal hypersurfaces 18
3.2. Construction of M-surfaces 19
4. Appendix 21
4.1. Usefull combinatorial lemmae 21
4.2. Formula for the number of k-simplices 23
References 25

The author is grateful to Max Planck Institut für Mathematik in Bonn for excellent working conditions. The author was partially supported by the European research network IHP-RAAG contract HPRN-CT-2001-00271.

INTRODUCTION

The number of connected components of a real algebraic plane projective curve of degree m is not more than $(m-1)(m-2)/2+1$. This bound was proved by Harnack who also showed that for any positive integer m there exist curves of degree m which are maximal in this sense (i.e. with $(m-1)(m-2)/2+1$ connected components). Harnack's bound is generalized to the case of any real algebraic variety by the Smith-Thom inequality. Let $b_i(V;K)$ be the i^{th} Betti number of a topological space V with coefficients in a field K (i.e. $b_i(V;K) = \dim_K(H_i(V;K))$). Denote by $b_*(V;K)$ the sum of the Betti numbers of V. Let X be a complex algebraic variety equipped with an anti-holomorphic involution c. The real part $\mathbb{R}X$ of X is the fixed point set of c. Then the Smith-Thom inequality states that $b_*(\mathbb{R}X;\mathbb{Z}_2) \leq b_*(X;\mathbb{Z}_2)$. A variety X for which $b_*(\mathbb{R}X;\mathbb{Z}_2) = b_*(X;\mathbb{Z}_2)$ is called a *maximal* variety or *M-variety.* The question "does a given family of real algebraic varieties contain maximal elements?" is one of the main problems in topology of real algebraic varieties. For the family of the hypersurfaces of a given degree in $\mathbb{R}P^d$ a positive answer is obtained in [IV05] using the combinatorial Viro method called T-construction (see [Vir84], [Vir04], [Ite97], and Theorem 1.1).

The Viro patchworking theorem –and its combinatorial particular case– is essential in the study of existence of hypersurfaces with prescribed topology. Let Δ be a lattice polytope equipped with a regular (convex) triangulation τ and a distribution of signs D at its vertices. From this data Viro's combinatorial patchwork produces a real algebraic hypersurface in the toric variety X_Δ associated to Δ whose topology can be easily recovered from Δ, τ and D (see the above references and Section 1.2). ***Primitive T-hypersurfaces*** are those obtained by the combinatorial patchworking if one starts with a unimodular (primitive) triangulation τ. Provided that Δ correspond to a nonsingular toric variety, we prove that for any primitive T-hypersurface Z,

$$\chi(\mathbb{R}Z) = \sigma(Z)\,, \tag{0.1}$$

where $\chi(\mathbb{R}Z)$ is the Euler characteristic of $\mathbb{R}Z$ and $\sigma(Z) := \sum_{p+q=0\,[2]}(-1)^p h^{p,q}(Z)$ is the signature of Z if $\dim Z$ is even and 0 otherwise (See Theorem 2.1).

This result can be naturally formulated in the tropical setting introduced by G. Mikhalkin. (See Section 1.3 or [Mik04] and [Mik05] for definitions and propreties of tropical hypersurfaces.):

The Euler characteristic of a compactified nonsingular real tropical hypersurface with Newton polytope Δ is equal to the signature of a complex algebraic hypersurface in X_Δ with Newton polytope Δ.

For primitive T-surfaces in the projective space $\mathbb{P}^3$ this result was first proven by I. Itenberg [Ite97] who used it to construct maximal T-surfaces in all degrees in $\mathbb{P}^3$. The existence of maximal surfaces in all degrees in the projective space had been proven in [Vir79] by Viro using small perturbation technics.

Using the combinatorial Viro method and equality (0.1) we prove the existence of maximal surfaces in a family of toric varieties which includes $\mathbb{P}^3$. A polytope Δ in $\mathbb{R}^d$ is a ***Nakajima polytope*** (see Figure 8) if either Δ is 0-dimensional or there exists a Nakajima polytope $\bar{\Delta}$ in $\mathbb{R}^{d-1}$ and a linear function $f : \mathbb{R}^{d-1} \to \mathbb{R}$, nonnegative on $\bar{\Delta}$, such that $f(\mathbb{Z}^{d-1}) \subset \mathbb{Z}$ and $\Delta = \{(x, x_d) \in \bar{\Delta} \times \mathbb{R} \mid 0 \leq x_d \leq f(x)\}$. We show that for any 3-dimensional Nakajima polytope Δ corresponding to a nonsingular toric variety there exists a maximal hypersurface in X_Δ with Newton polytope Δ.

I am grateful to Ilia Itenberg for his valuable advice and to Erwan Brugallé for his pertinent remarks.

Organization of the material. We first introduce notation and describe combinatorial patchworking in Section 1. In the same section, we describe briefly tropical hypersurfaces and state the results in the tropical setting. We also recall there facts about lattice triangulations and state Danilov and Khovanskii Formulae. Section 2 is devoted to the proof of equality (0.1) for primitive hypersurfaces stated in Theorem 2.1. In the third section we construct maximal surfaces in a family of toric varieties using equality (0.1). We postpone some proofs of Lemmae needed in Section 2 to the appendix (Section 4) where we also give some usefull combinatorial formulae (e.g. Lemma 6).

1. Preliminaries

1.1. **Toric varieties.** We fix here some conventions and notations, the construction of toric varieties we use is based on the one described in [Ful93]. Fix an orthonormal basis of $\mathbb{R}^d$ and thus an inclusion $\mathbb{Z}^d \to \mathbb{R}^d$. This inclusion defines a lattice N in $\mathbb{R}^d$. Denote by M the dual lattice $\mathrm{Hom}(N, \mathbb{Z})$ of N. Here we consider polytopes in $M \otimes \mathbb{R}$. By ***polytope*** we mean convex polytope whose vertices are integer (i.e. belong to the lattice M). Let Δ be a polytope in $M \otimes \mathbb{R}$. Let p be a vertex of Δ and let $\Gamma_1, \cdots, \Gamma_k$ be the facets of Δ containing p. To p we associate the cone σ_p generated by the minimal integer inner normal vectors of $\Gamma_1, \cdots, \Gamma_k$. The inner normal fan $\mathfrak{E}_\Delta$ is the fan whose d-dimensional cones are the cones σ_p for all vertices p of Δ. The toric variety X_Δ associated to Δ is the toric variety $X(\mathfrak{E}_\Delta)$ associated to the fan $\mathfrak{E}_\Delta$.

1.2. **Combinatorial patchworking.** Let us briefly describe the *combinatorial patchworking*, also called *T-construction* , which is a particular case of the Viro method. A more detailed exposition can be found in [IV05], [Vir04] or [GKZ94] p. 385.

By a ***subdivision*** of a polytope we mean a subdivision in convex polytopes whose vertices have integer coordinates. A subdivision τ of a d-dimensional polytope Δ is called ***convex*** (or ***regular***) if there exists a convex piecewise-linear function $\Phi : \Delta \to \mathbb{R}$ whose domains of linearity coincide with the d-dimensional polytopes of τ. A triangulation is a subdivision into simplices.

Given a triple (Δ, τ, D), where Δ is a polytope, τ a convex triangulation of Δ, and D a distribution of signs at the vertices of τ (i.e. each vertex is labelled with $+$ or $-$), the combinatorial patchworking, produces an algebraic hypersurface Z in X_Δ.

Let Δ be a d-dimensional polytope in the positive orthant $(\mathbb{R}^+)^d = \{(x_1, \dots x_d) \in \mathbb{R}^d \mid x_1 \geq 0, \dots, x_d \geq 0\}$, and τ be a convex triangulation of Δ. Denote by $s_{(i)}$ the reflection with respect to the coordinate hyperplane $x_i = 0$ in $\mathbb{R}^d$. Consider the union Δ^* of all copies of Δ under the compositions of reflections $s_{(i)}$ and extend τ to a triangulation τ^* of Δ^* by means of these reflections. We extend $D(\tau)$ to a distribution of signs at the vertices of τ^* using the following rule : for a vertex a of τ^*, one has $\mathrm{sign}(s_{(i)}(a)) = \mathrm{sign}(a)$ if the i-th coordinate of a is even, and $\mathrm{sign}(s_{(i)}(a)) = -\mathrm{sign}(a)$, otherwise.

Let σ be a d-dimensional simplex of τ^* with vertices of different signs, and E be the piece of hyperplane which is the convex hull of the middle points of the edges of σ with endpoints of opposite signs. We separate vertices of σ labelled with $+$ from vertices labelled with $-$ by E. The union of all these hyperplane pieces forms a piecewise-linear hypersurface H^*.

For any facet Γ of Δ^*, let N^Γ be a vector normal to Γ. Let F be a face of Δ^* and $\Gamma_1, \dots, \Gamma_k$ be the facets containing F. Let L be the linear space spanned by $N^{\Gamma_1}, \dots, N^{\Gamma_k}$. For any $v = (v_1, \dots, v_d) \in L \cap \mathbb{Z}^d$ identify F with ${s_{(1)}}^{v_1} \circ {s_{(2)}}^{v_2} \circ \cdots \circ {s_{(d)}}^{v_d}(F)$. Denote by $\widetilde{\Delta}$ the result of the identifications. The variety $\widetilde{\Delta}$ is homeomorphic to the real part $\mathbb{R}X_\Delta$ of X_Δ (see, for example, [GKZ94] Theorem 5.4 p. 383 or [Stu94] Proposition 2).

Denote by $\widetilde{H}$ the image of H^* in $\widetilde{\Delta}$. Let Q be a polynomial with Newton polytope Δ. It defines a hypersurface Z_0 in the torus $(\mathbb{C}^*)^d$ contained in X_Δ. The closure Z of Z_0 in X_Δ is the hypersurface defined by Q in X_Δ. We call Δ the ***Newton polytope*** of Z.

Theorem 1.1 (T-construction, O. Viro)**.** *Under the hypotheses made above, there exists a hypersurface Z of X_Δ with Newton polytope Δ and a homeomorphism $h : \mathbb{R}X_\Delta \to \widetilde{\Delta}$ such that $h(\mathbb{R}Z) = \widetilde{H}$.*

A ***real algebraic T-hypersurface*** is a hypersurface which can be obtained via the above construction. A d-dimensional simplex with integer vertices is called ***primitive*** (or ***unimodular***) if its volume is equal to $\frac{1}{d!}$. A triangulation τ of a d-dimensional polytope is ***primitive*** if every d-simplex of the triangulation is primitive. A T-hypersurface Z is ***primitive*** if it can be constructed with the combinatorial patchworking using a primitive triangulation.

Let Δ be a d-dimensional polytope and τ a (convex) triangulation of Δ. The star, $\mathrm{Star}(v)$, of a vertex v of τ is the union of all simplices of τ containing v. Suppose that τ is equipped with a distribution of signs. We say that v is ***isolated*** if all other vertices of $\mathrm{Star}(v)$ have the sign opposite to $\mathrm{sign}(v)$.

Remark 1. *Let p be an isolated vertex of the triangulation τ^* of Δ^*. Assume that p is in the interior of Δ^*. Then the star of p contains a connected component of H^* homeomorphic to a sphere S^{d-1}. So, p corresponds to a $(d-1)$-dimensional sphere of the real part of the hypersurface Z obtained by Viro's theorem.*

Notice that the triangulation τ^* induces a natural cell decomposition of H^*: each cell is the intersection of a simplex of τ^* with H^*. A k-dimensional simplex in $\mathbb{R}^d$ is called ***elementary*** if the reductions modulo 2 of its vertices generate a k-dimensional affine space over $\mathbb{Z}_2$.

Our computation of the Euler characteristic of $\mathbb{R}Z$ uses the following proposition due to Itenberg (see [Ite97] Proposition 3.1).

Proposition 1.2. *Let s be an elementary k-simplex of a triangulation τ of a d-dimensional polytope. Assume that s is contained in j coordinate hyperplanes. Then the union s^* of the symmetric copies of s contains exactly $(2^d - 2^{d-k})/2^j$ cells of dimension $k-1$ of the cell decomposition of H^*.*

1.3. **Tropical hypersurfaces.** Before stating the results we briefly recall some basic facts about tropical hypersurfaces. A detailed presentation can be found in [Mik05]. In this article we will use the definition of tropical varieties as images of algebraic varieties defined over the field of Puiseux series.

Let $\mathbb{K}$ be the field of Puiseux series. An element of $\mathbb{K}$ is a series $g(t) = \sum_{r \in R} b_r t^r$ where $b_r \in \mathbb{C}$, $R \subset \mathbb{Q}$ is bounded from below and contained in an arithmetic sequence. Consider the valuation $\operatorname{val}(g(t)) := \min\{r | b_r \neq 0\}$. For convenience we choose Mikhalkin's conventions and consider in fact minus the valuation, $v(g) := -\operatorname{val}(g)$. Let f be a polynomial in $\mathbb{K}[z_1, \cdots, z_n] = \mathbb{K}[z]$. It is of the form $f(z) = \sum_{\omega \in A} c_\omega z^\omega$ with A a finite subset of $\mathbb{Z}^n$ and $c_\omega \in \mathbb{K}^*$. Consider $Z_f := \{z \in (\mathbb{K}^*)^n | f(z) = 0\}$ the zero set of f in $(\mathbb{K}^*)^n$ and the coordinatewise valuation (up to sign)

$$\begin{array}{rccl} V: & (\mathbb{K}^*)^n & \longrightarrow & \mathbb{R}^n \\ & z & \longmapsto & (v(z_1), \ldots, v(z_n)). \end{array}$$

Definition 1.1. *The tropical hypersurface Z_f^{trop} associated to f is the closure (in the usual topology) of the image under V of the hypersurface Z_f:*

$$Z_f^{\text{trop}} := \overline{V(Z_f)} \subset \mathbb{R}^n.$$

The following theorem, due to Kapranov, allows one to see tropical hypersurfaces as nonlinearity domains of some piecewise-linear convex functions. Put

$$\begin{array}{rccl} \nu: & A & \longrightarrow & \mathbb{R} \\ & \omega & \longmapsto & -v(c_\omega)\,, \end{array}$$

the Legendre transform $\mathcal{L}(\nu)$ of ν is the piecewise-linear convex function defined as follows:

$$\begin{array}{rccl} \mathcal{L}(\nu): & \mathbb{R}^n & \longrightarrow & \mathbb{R} \\ & x & \longmapsto & \max_{\omega \in A}(x \cdot \omega - \nu(\omega)) \end{array}$$

Theorem 1.2 (Kapranov)**.** *The tropical hypersurface Z_f^{trop} is the nonlinearity domain of $\mathcal{L}(\nu)$:*

$$\overline{V(Z_f)} = \textit{corner locus}(x \mapsto \max_{\omega \in A}(x \cdot \omega + v(c_\omega))).$$

Another very nice fact is that one can naturally associate to a tropical hypersurface a dual subdivision of its Newton polytope. Let Γ be the convex hull in $\mathbb{R}^n \times \mathbb{R}$ of the points $(\omega, v(c_\omega))$ for all ω in A. Put

$$\begin{array}{rccl} \overline{\nu}: & \Delta & \longrightarrow & \mathbb{R} \\ & x & \longmapsto & \min\{y | (x, y) \in \Gamma\}. \end{array}$$

The linearity domains of $\overline{\nu}$ are the n-cells of a convex polyhedral subdivision τ of Δ. The hypersurface Z_f^{trop} induces a subdivision Ξ of $\mathbb{R}^n$. The subdivisions τ and Ξ are dual in the following sense (see Figure 1).

There is a one-to-one inclusion reversing correspondence L between cells of Ξ and cells of τ such that for any $\xi \in \Xi$,



FIGURE 1. A tropical line, a tropical cubic with the dual subdivision of its Newton polytope.

(1) $\dim L(\xi) = \operatorname{codim} \xi$,
(2) the affine supports of $L(\xi)$ and ξ are orthogonal.

We say that Z_f^{trop} is **nonsingular** if its dual subdivision τ is a primitive triangulation.

Complex tropical hypersurfaces. In order to prove his correspondence theorem Mikhalkin introduces in [Mik05] the complexification of tropical curves. Let $g(t) = \sum_{r \in R} b_r t^r$ be an element of $\mathbb{K}^*$. Define the argument $\arg(g)$ to be the argument of the coefficient of the smallest power of t appearing in g (i.e. $\arg(g) = \arg(b_{\text{val}(g)})$). Let Arg be the coordinatewise argument on $(\mathbb{K}^*)^n$. Consider the "complexification" of the valuation

$$\begin{array}{rrcl} W := V \times \operatorname{Arg}: & (\mathbb{K}^*)^n & \longrightarrow & \mathbb{R}^n \times (S^1)^n \\ & z & \longmapsto & ((v(z_1), \ldots, v(z_n)), (\arg(z_1), \ldots, \arg(z_n))). \end{array}$$

One can define complex tropical hypersurfaces to be the closure of the image under W of hypersurfaces in $(\mathbb{K}^*)^n$ but it is also convenient to consider the composition of W with the exponential. Namely, put

$$\begin{array}{rrcl} V_{\mathbb{C}}: & (\mathbb{K}^*)^n & \longrightarrow & (\mathbb{C}^*)^n \\ & z & \longmapsto & (e^{v(z_1) + i \arg(z_1)}, \ldots, e^{v(z_n) + i \arg(z_n)}) \end{array}$$

We will call both homeomorphic objects complex tropical hypersurfaces and use one and the other alternatively depending on the context.

Definition 1.3. *The complex tropical hypersurface* $\mathbb{C}Z_{f,V_{\mathbb{C}}}^{\text{trop}}$ *(resp.* $\mathbb{C}Z_{f,W}^{\text{trop}}$*) associated to* f *is the closure of the image under* $V_{\mathbb{C}}$ *(resp.* W*) of the hypersurface* Z_f*:*

$$\mathbb{C}Z_{f,V_{\mathbb{C}}}^{\text{trop}} := \overline{V_{\mathbb{C}}(Z_f)} \subset (\mathbb{C}^*)^n,$$

or alternatively

$$\mathbb{C}Z_{f,W}^{\text{trop}} := \overline{W(Z_f)} \subset \mathbb{R}^n \times (S^1)^n.$$



FIGURE 2. A complex tropical line, its tropical "projection" and a real tropical line in the four quadrants.

For example, a complex tropical line is homeomorphic to a sphere with three punctures (see Figure 2).

Real tropical Hypersurfaces. Assume from now on that $f = \sum_{\omega \in A} c_\omega z^\omega$ is real (i.e. all the coefficients a_r of each series $c_\omega = \sum_{r \in R} a_r t^r$ are real).

Definition 1.4. *The real tropical hypersurface* $\mathbb{R}Z^{\text{trop}}_{f,V_\mathbb{C}}$ *associated to* f *is the intersection in* $(\mathbb{C}^*)^n$ *of* $\mathbb{C}Z^{\text{trop}}_{f,V_\mathbb{C}}$ *with* $(\mathbb{R}^*)^n$*:*

$$\mathbb{R}Z^{\text{trop}}_{f,V_\mathbb{C}} := \overline{V_\mathbb{C}(Z_f)} \cap (\mathbb{R}^*)^n,$$

or alternatively

$$\mathbb{R}Z^{\text{trop}}_{f,W} := \overline{W(Z_f)} \cap (\mathbb{R}^n \times \{0, \pi\}^n).$$

On Figure 2 we have pictured the intersection of a real tropical line with each of the four quadrants. The sign $\operatorname{sign}(c_\omega)$ of a coefficient $c_\omega = \sum_{r \in R} a_r t^r$ of f is defined to be the sign of the first coefficient $a_{\operatorname{val}(c_\omega)}$ of c_ω (i.e. $\operatorname{sign} c_\omega := \operatorname{sign} a_{\operatorname{val}(c_\omega)}$). From now on we will only consider nonsingular tropical hypersurfaces. For $\epsilon \in \{+1, -1\}^n$, let $\mathcal{Q}_\epsilon$ be the orthant of $(\mathbb{R}^*)^n$ which maps to the positive orthant $(\mathbb{R}^*_+)^n$ under the isometry $\varphi_\epsilon : \mathcal{Q}_\epsilon \to (\mathbb{R}^*_+)^n$ defined by $\varphi_\epsilon(x_1, \cdots, x_n) = (\epsilon_1 x_1, \cdots, \epsilon_n x_n)$. If $Z^{\text{trop}}_{f,V_\mathbb{C}}$ is nonsingular one can reconstruct $\mathbb{R}Z^{\text{trop}}_f \cap \mathcal{Q}_\epsilon$ only from the data of Z^{trop}_f and the collection of signs $\operatorname{sign} c_\omega$ of the coefficients of f (see [Mik04] pp. 25 and 37, [Vir01], and [Mik00] Appendix for the case of amoebas). First one sees that we can restrict the study to the case of the first orthant $\mathcal{Q}$ since one can map any $\mathcal{Q}_\epsilon$ to $\mathcal{Q}$ by φ_ϵ and change f to $f_\epsilon = f(\epsilon_1 x_1, \cdots, \epsilon_n x_n)$.

Let us rather use the map W (it suffices to exponentiate to switch back to the image under $V_\mathbb{C}$). Then $\mathcal{Q}$ corresponds to $\mathbb{R}^n \times \{(0, \cdots, 0)\}$. Consider the tropical hypersurface $Z^{\text{trop}}_f \subset \mathbb{R}^n$, the induced subdivision Ξ_f of $\mathbb{R}^n$ and the dual subdivision τ_f of its Newton polytope Δ_f. Let D_f be the sign distribution at the vertices of τ_f such that a vertex ω is labelled with the sign $\operatorname{sign}(c_\omega)$ of the corresponding coefficient in f.

Let x be a point of Z^{trop}_f. The point $(x, (0, \cdots, 0))$ is in $\mathbb{R}Z^{\text{trop}}_{f,W} \cap (\mathbb{R}^n \times \{(0, \cdots, 0)\})$ if and only if x belongs to the closure of an $(n-1)$-cell ξ of Ξ_f which is dual to an edge



FIGURE 3. A real tropical cubic in the first quadrant, the underlying tropical curve (in dotted lines) and its dual subdivision.



FIGURE 4. An attempt at representing a real nonsingular tropical cubic with the underlying symmetrized tropical curve (in dotted lines) and corresponding T-curve.

of τ_f whose extremities have opposite signs (see Figure 3 for an example in the case of a curve).

Then one sees that one can associate a unique primitive T-hypersurface to a nonsingular real tropical hypersurface. Namely take the T-hypersurface H constructed from Δ_f, τ_f and D_f. Clearly, there exists a homeomorphism $h : (\mathbb{R}^*)^n \to (\Delta \setminus \partial\Delta)^*$ such that $h(\mathbb{R}Z_f^{\text{trop}}) = H \cap (\Delta \setminus \partial\Delta)^*$. On Figure 4 we tried to illustrate this homeomorphism in the case of curves.

The results in this paper are nicer and much easier to state when one compactifies the tropical hypersurfaces. Let us describe the natural compactification of a tropical hypersurface. Recall that $\mathbb{C}Z_{f,V_{\mathbb{C}}}^{\text{trop}} = \overline{V_{\mathbb{C}}(Z_f)}$ is a subset of the torus $(\mathbb{C}^*)^n$. Consider the usual compactification of $(\mathbb{C}^*)^n$ into the toric variety X_{Δ_f} associated to the Newton polytope Δ_f of f. Let $\iota : (\mathbb{C}^*)^n \hookrightarrow X_{\Delta_f}$ be the corresponding inclusion. We define the compactification $\overline{\mathbb{C}Z_f^{\text{trop}}}$ to be the closure of $\iota(\mathbb{C}Z_{f,V_{\mathbb{C}}}^{\text{trop}})$ in X_{Δ_f}. Note that the stratification of X_{Δ_f} into orbits of the action of $(\mathbb{C}^*)^n$ defines a natural stratification of $\overline{\mathbb{C}Z_f^{\text{trop}}}$.

We sum up natural maps in the following commutative diagram.

$$\begin{tikzcd}
\mathbb{R}Z^{\text{trop}}_{f,W} \arrow[r, "\sim"] \arrow[d, hook] & \mathbb{R}Z^{\text{trop}}_{f,V_{\mathbb{C}}} \arrow[r, hook] \arrow[d, hook] & (\mathbb{R}^*)^n \arrow[r, hook, "\iota_{\mathbb{R}}"] \arrow[d, hook] & \mathbb{R}X_{\Delta_f} \arrow[d, hook] \\
\mathbb{C}Z^{\text{trop}}_{f,W} \arrow[r, "\sim"] & \mathbb{C}Z^{\text{trop}}_{f,V_{\mathbb{C}}} \arrow[r, hook] & (\mathbb{C}^*)^n \arrow[r, hook, "\iota"] & X_{\Delta_f}
\end{tikzcd}$$

Define $\overline{\mathbb{R}Z^{\text{trop}}_f}$ to be the intersection of $\overline{\mathbb{C}Z^{\text{trop}}_f}$ with the real part $\mathbb{R}X_{\Delta_f}$ of X_{Δ_f}. Clearly $\overline{\mathbb{R}Z^{\text{trop}}_f}$ is also the closure of $\iota_{\mathbb{R}}(\mathbb{R}Z^{\text{trop}}_f)$ in $\mathbb{R}X_{\Delta_f}$. One can see that the natural stratification of $\overline{\mathbb{R}Z^{\text{trop}}_f}$ induced by the torus action corresponds to the stratification of the T-hypersurface H induced by the face complex of Δ_f which will be used in the proof of Theorem 2.1.

1.4. **Tropical Statements.** We can now state our results in the tropical language. Let Δ be a polytope corresponding to a nonsingular toric variety.

Theorem 1.3. *Let $\overline{\mathbb{R}Z^{\text{trop}}_f}$ be a compactified nonsingular real tropical hypersurface with Newton polytope $\Delta_f = \Delta$ then*

$$\chi(\overline{\mathbb{R}Z^{\text{trop}}_f}) = \sigma(Z),$$

where Z is a smooth complex algebraic hypersurface in X_{Δ_f} with Newton polytope Δ_f and $\sigma(Z) = \sum_{p+q=0\,[2]} (-1)^p h^{p,q}(Z)$.

Theorem 1.3 is equivalent to Theorem 2.1 about Euler characteristic of primitive T-hypersurfaces and is one the of main results of this paper.

One can define a maximal real tropical hypersurface in the following way. Let f be a polynomial over $\mathbb{K}$ such that every coefficient of f is a series with only real coefficients. Let Δ_f be the Newton polytope of f. The tropical hypersurface $\overline{\mathbb{R}Z^{\text{trop}}_f}$ is **maximal** if

$$b_*(\overline{\mathbb{R}Z^{\text{trop}}_f}; \mathbb{Z}_2) = b_*(\overline{\mathbb{C}Z^{\text{trop}}_f}; \mathbb{Z}_2).$$

We can then prove existence results for maximal real tropical surfaces.

Theorem 1.4. *Let Δ be a 3-dimensional Nakajima polytope corresponding to a nonsingular toric variety X_Δ. Then, there exists a maximal real tropical surface in X_Δ with the Newton polytope Δ.*

Theorem1.4 will be proved in Section 3 (See Theorem 3.3) as well as the following corollary (See Corollary 3.4).

For a nonnegative integer α and positive integers m and n denote by $\delta^{m,n}_\alpha$ the polygon having the vertices $(0,0)$, $(n+m\alpha, 0)$, $(0,m)$, and (n,m) in $\mathbb{R}^2$. The toric variety associated with $\delta^{m,n}_\alpha$ is a rational ruled surface Σ_α. Consider now the truncated cylinder $\Delta^{l,m,n}_\alpha$ with base $\delta^{m,n}_\alpha$ whose vertices are

$$(0,0,0), (n+m\alpha,0,0), (0,m,0), (n,m,0),$$
$$(0,0,l), (n+m\alpha,0,l), (0,m,l), \text{ and } (n,m,l),$$

where l is a positive integer. The toric variety $X_{\Delta_\alpha^{l,m,n}}$ associated with $\Delta_\alpha^{l,m,n}$ is $\Sigma_\alpha \times \mathbb{C}P^1$. Since $\Delta_\alpha^{l,m,n}$ is, up to exchanging two coordinates, a Nakajima polytope, we have the foolowing corollary (See Corollary 3.4).



FIGURE 5. Polygon $\delta_\alpha^{m,n}$.



FIGURE 6. Polytope $\Delta_\alpha^{l,m,n}$.

Corollary 1.5. *For any nonnegative integer α and any positive integers m, n and l, there exists a (real) maximal nonsingular tropical surface $\overline{\mathbb{R}Z_f^{\text{trop}}}$ with the Newton polytope $\Delta_\alpha^{l,m,n}$ in $\mathbb{R}\Sigma_\alpha \times \mathbb{R}P^1$.*

1.5. **Some facts about triangulations of lattice polytopes.**

1.5.1. *Ehrhart Polynomial.* In order to compute both sides of equality (0.1), namely the Euler characteristic of the real part of a primitive T-hypersurface and the signature of its complex part, we use some combinatorics of triangulations of lattice polytopes. We first recall the results of E. Ehrhart on the number of integer points in a lattice polytope. Ehrhart showed that the number of integer points in the multiple $\lambda \cdot \Delta$ of a (not necessarily convex) polytope Δ, where λ is a positive integer, is a polynomial in λ (see [Ehr94] or [Ehr67]). We denote by $l(\Delta)$ and $l^*(\Delta)$ the numbers of integer points in Δ and in the interior of Δ, respectively.

Theorem 1.6 (E. Ehrhart)**.** *Let* Δ *be a polytope with integer vertices. Then, the numbers* $l(\lambda \cdot \Delta)$ *and* $l^*(\lambda \cdot \Delta)$ *are polynomials in* λ *of degree* $\dim \Delta$*. Denote them respectively by* $Ehr_\Delta(\lambda)$ *and* $Ehr^*_\Delta(\lambda)$*. They satisfy the reciprocity law*

$$(-1)^{dim\Delta} Ehr_\Delta(-\lambda) = Ehr^*_\Delta(\lambda).$$

One often considers the Ehrhart series

$$SE_p(t) = \sum_{\lambda=0}^{\infty} Ehr_\Delta(\lambda) t^\lambda.$$

Put $Q_\Delta(t) = (1-t)^{d+1} SE_p(t)$, where d is the dimension of Δ. We define the numbers Ψ_j to be the coefficients of $Q_\Delta(t)$:

$$Q_\Delta(t) = \sum_{j=0}^{\infty} \Psi_j t^j.$$

In fact, $Q_\Delta(t)$ is a polynomial of degree d (see [Bri94] or lemma 1). Let a_i^Δ be the coefficient of λ^i in $Ehr_\Delta(\lambda)$. The following formula can be found in Section 4.1 of [DHZ01] (See also the Appendix).

Lemma 1. *One has*

$$\Psi_j = \sum_{i=0}^{d} (\sum_{n=0}^{j} (-1)^{j-n} \binom{d+1}{j-n} n^i) a_i^\Delta,$$

and $\Psi_j = 0$ *for* $j \geq d+1$.

1.5.2. *Number of simplices in a primitive triangulation.* Another usefull formula expresses the numbers of simplices of any dimension in the primitive triangulation τ of Δ in terms of the coefficients of the Ehrhart polynomial of Δ. In the case of a primitive triangulation, these numbers happen not to depend on the primitive triangulation chosen. This statement can be found in [Dai00] and in Appendix 4.2 we provide a proof in the case of convex triangulations.

Let nbs_r^F be the number of r-dimensional simplices in a primitive triangulation of F which are contained in the interior of F, and let i be the dimension of F. Let S_2 be the Stirling number of the second kind defined by $S_2(i,j) = 1/(j)! \sum_{k=0}^{j} (-1)^{j-k} \binom{j}{k} k^i$.

Proposition 1.7. *Under the above hypotheses we have,*

$$N_r^F = \sum_{l=r+1}^{i+1} (-1)^{i-l+1} \cdot r! S_2(l, r+1) \cdot a_{l-1}^F.$$

1.6. **Danilov and Khovanskii formulae.** V. Danilov and A. Khovanskii [DK87] computed the Hodge numbers of a smooth hypersurface in a toric variety X_Δ in terms of the polytope Δ. Recall that, for a face F of Δ, the coefficient of the term of degree l of the Ehrhart polynomial of F is denoted by a_l^F. Let $\mathcal{F}_i(\Delta)$ be the set of i-dimensional faces of Δ, and $f_i(\Delta)$ be the cardinality of $\mathcal{F}_i(\Delta)$. A d-dimensional ***simple*** polytope is one for which each vertex is contained in exactly d edges.

Theorem 1.8 (V. Danilov, A. Khovanskii)**.** *Let Δ be a simple polytope of dimension d, and Z be a smooth or quasi-smooth*[1] *algebraic hypersurface with Newton polytope Δ in X_Δ. Then, for $p \neq \frac{d-1}{2}$*

$$\begin{aligned} h^{p,p}(Z) &= (-1)^{p+1}\sum_{i=p+1}^{d}(-1)^i\binom{i}{p+1}f_i(\Delta) \\ h^{\frac{d-1}{2},\frac{d-1}{2}}(Z) &= (-1)^{\frac{d+1}{2}}\sum_{i=\frac{d+1}{2}}^{d}(-1)^i\binom{i}{\frac{d+1}{2}}f_i(\Delta) - \sum_{i=\frac{d+1}{2}}^{d}\sum_{F\in\mathcal{F}_i(\Delta)}(-1)^i\Psi_{\frac{d+1}{2}}(F) \\ h^{p,d-1-p}(Z) &= (-1)^d\sum_{i=p+1}^{d}\sum_{F\in\mathcal{F}_i(\Delta)}(-1)^i\Psi_{p+1}(F) \\ h^{p,q} &= 0 \text{ if } q \neq p \text{ or } p \neq d-1-p, \end{aligned}$$

where $\Psi_{p+1}(F) = \sum_{\alpha=1}^{i+1}\sum_{a=0}^{p+1}(-1)^a\binom{i+1}{a}(p+1-a)^{\alpha-1}a^F_{\alpha-1}$.

2. PRIMITIVE T-HYPERSURFACES

2.1. **Statement.** Let Δ be a d-dimensional polytope corresponding to a nonsingular toric variety. Let Z be a primitive T-hypersurface with Newton polytope Δ in X_Δ, and τ be a primitive triangulation from which one can construct Z via the combinatorial Viro method. Notice that since X_Δ is smooth the T-hypersurface Z is also smooth. Define $\sigma(Z)$ by

$$\sigma(Z) := \sum_{p+q=0\ [2]}(-1)^p h^{p,q}(Z).$$

If d is odd and Z is smooth, the intersection form

$$\iota : H_{d-1}(Z;\mathbb{Z})/tors \times H_{d-1}(Z;\mathbb{Z})/tors \to \mathbb{Z}$$

is a symmetric bilinear form and its signature is called the signature of Z and is equal to $\sigma(Z)$. (See, for example, [GH78] Chapter 0, Section 7 p. 126). We have the following statement.

Theorem 2.1. *If Z is a primitive real algebraic T-hypersurface in a nonsingular toric variety X_Δ, then*

$$\chi(\mathbb{R}Z) = \sigma(Z),$$

where $\chi(\mathbb{R}Z)$ is the Euler characteristic of the real part $\mathbb{R}Z$ of Z.

Remark 2.1. For simplicity we state Theorem 2.1 for a nonsingular ambiant toric variety. In fact, we only need to require that the polytope Δ be simple (see Subsection 1.6). In this case X_Δ is quasi-smooth (i.e. locally isomorphic to a toric variety defined by a simplicial cone; see [Dan78] Section 14). Then Z is also quasi-smooth (see [Dan78] Section 13), the Hodge numbers can be calculated by Danilov and Khovanskii formulae (see Theorem 1.8) and the proof is verbatim.

---

[1] See [Dan78] and Remark 2.1

Remark 2.2. The signature $\sigma(Z)$ is the same for any generic hypersurface Z with Newton polytope Δ in X_Δ but if Z is not a primitive T-hypersurface $\chi(\mathbb{R}Z)$ needs not be equal to the signature (and very often is not).

Recall that $l^*(\Delta)$ is the number of integer points in the interior of Δ. Here and further on we write $b_i(X)$ for $b_i(X;\mathbb{Z}_2)$.

Corollary 2.2. *Assume that Δ is a 3-dimensional simple polytope and Z is a primitive T-surface in X_Δ with Newton polytope Δ. If the number $b_0(\mathbb{R}Z)$ of connected components of $\mathbb{R}Z$ is at least $l^*(\Delta)+1$, then $b_0(\mathbb{R}Z) = l^*(\Delta)+1$ and Z is maximal.*

Proof. First, note that $l^*(\Delta) = h^{2,0}(Z)$. Then $\chi(\mathbb{R}Z) \geq 2h^{2,0}(Z)+2-b_1(\mathbb{R}Z)$. Now using the equalities $\chi(\mathbb{R}Z) = \sigma(Z) = 2h^{2,0}(Z)+2-h^{1,1}(Z)$ one gets $b_1(\mathbb{R}Z) \geq h^{1,1}(Z)$. Furthermore $h^{1,0}(Z) = h^{0,1}(Z) = 0$ (see Theorem 1.8), and thus $b_*(Z;\mathbb{C}) = 2h^{2,0}(Z)+2+h^{1,1}(Z)$. Hence $b_*(\mathbb{R}Z;\mathbb{Z}_2) \geq b_*(Z;\mathbb{C}) = b_*(Z;\mathbb{Z}_2)$. The Smith-Thom inequality implies that $b_*(\mathbb{R}Z;\mathbb{Z}_2) = b_*(Z;\mathbb{Z}_2)$, and thus, that $b_1(\mathbb{R}Z) = h^{1,1}(Z)$ and $b_0(\mathbb{R}Z) = h^{2,0}(Z)+1$. □

2.2. **Proof of Theorem 2.1.** If d is even, Theorem 2.1 is straightforward. Indeed, in this case Z is a (smooth) odd dimensional hypersurface, so $\chi(\mathbb{R}Z) = 0$. On the other hand, we have the equality $h^{p,q}(Z) = h^{d-1-p,d-1-q}(Z)$ for any p and q, and $d-1$ is odd. Thus, $\sigma(Z) = \sum_{p+q=0\,[2]}(-1)^p h^{p,q}(Z) = 0$.

Assume now that d is odd.

Lemma 2. *We have*

$$\chi(\mathbb{R}Z) = \sum_{i=1}^{d}\sum_{F\in\mathcal{F}_i(\Delta)}\sum_{l=2}^{i+1}\chi_{l,i+1}a^F_{l-1} \quad \textit{and} \quad \sigma(Z) = \sum_{i=1}^{d}\sum_{F\in\mathcal{F}_i(\Delta)}\sum_{l=2}^{i+1}\sigma_{l,i+1}a^F_{l-1},$$

where

$$\begin{aligned} \chi_{l,i+1} &= (-1)^{i-l+1}\sum_{k=1}^{i}\frac{(2^i-2^{i-k})}{k+1}\sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l, \\ \sigma_{l,i+1} &= \sum_{p=0}^{d-1}(-1)^i\sum_{q=0}^{p+1}(-1)^{p+1-q}\binom{i+1}{q}(p+1-q)^{l-1}. \end{aligned}$$

Proof. The triangulation τ^* induces a cell decomposition $\mathcal{D}$ of $\widetilde{H}$. Let $\widetilde{I_F}$ be the image in $\widetilde{\Delta}$ of the union of the symmetric copies of the interior of a face F and $\mathcal{D}(\widetilde{I_F})$ be the set of cells of $\mathcal{D}$ contained in $\widetilde{I_F}$. Put $\chi_F = \sum_{\delta\in\mathcal{D}(\widetilde{I_F})}(-1)^{dim(\delta)}$. Then $\chi(\mathbb{R}Z) = \sum_{i=1}^{d}\sum_{F\in\mathcal{F}_i(\Delta)}\chi_F$. According to Proposition 1.2, if Q is a k-simplex of τ contained in j coordinate hyperplanes then the union of the symmetric copies of Q contains exactly $(2^d-2^{d-k})/2^j$ cells of dimension $k-1$. An i-face F of Δ^* contained in j coordinate hyperplanes is identified with $2^{d-i-j}-1$ other copies of F when passing from Δ^* to $\widetilde{\Delta}$. Thus, the number of

$(k-1)$-cells in $\mathcal{D}(\widetilde{I_F})$ is equal to $\frac{(2^d-2^{d-k})}{2^{d-i}}N_k^F$, where N_k^F is the number of k-simplices in the interior of F. Thus,

$$\chi_F = \sum_{k=1}^{\dim F} (-1)^{k-1}(2^{\dim F} - 2^{\dim F - k})N_k^F \ . \tag{2.1}$$

According to Proposition 1.7 of Section 1.5.2

$$N_k^F = \sum_{l=k+1}^{\dim F+1} k! S_2(l, k+1)(-1)^{\dim F - l+1} a_{l-1}^F,$$

where $S_2(i,j) = 1/j! \sum_{k=0}^{j}(-1)^{j-k}\binom{j}{k}k^i$ is the Stirling number of the second kind. Then one has

$$\begin{aligned}
\chi_F &= \sum_{k=1}^{\dim F} (-1)^{k-1}(2^{\dim F} - 2^{\dim F-k})N_k^F \\
&= \sum_{k=1}^{\dim F} (-1)^{\dim F-1}\frac{2^{\dim F} - 2^{\dim F-k}}{k+1}\sum_{l=k+1}^{\dim F+1}(-1)^l a_{l-1}^F \sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l \ .
\end{aligned}$$

By Lemma 6, Equality (4.2) (see Appendix), the sum on l can be taken from $l=2$ instead of $l=k+1$ and one gets

$$\chi_F = \sum_{l=2}^{\dim F+1}(-1)^l a_{l-1}^F \sum_{k=1}^{\dim F}(-1)^{\dim F-1}\frac{2^{\dim F} - 2^{\dim F-k}}{k+1}\sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l \ ,$$

which finishes the proof of the formula for $\chi(\mathbb{R}Z)$.

To compute $\sigma(Z)$, we use the Danilov and Khovanskii formulae (see Theorem 1.8). One obtains the following expression for $\sigma(Z)$:

$$\sigma(Z) \ = \ \sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i \sum_{F\in\mathcal{F}_i(\Delta)}\left(-\binom{i}{p+1} + (-1)^{p+1}\sum_{l=1}^{i+1}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1}a_{l-1}^F\right).$$

Consider $\sigma(Z)$ as an affine polynomial in the variables a_l^F. Denote by σ^{cst} the constant term of $\sigma(Z)$, and by σ^1 the sum of monomials in variables a_0^F. We have

$$\sigma^{cst} = -\sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i\binom{i}{p+1}f_i(\Delta)$$

and

$$\sigma^1 = \sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i(-1)^{p+1}\sum_{F\in\mathcal{F}_i(\Delta)}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}a_0^F.$$

Lemma 3. *One has* $\sigma^{cst} + \sigma^1 = 0$.

Proof. Since $a_F^0 = 1$, we obtain

$$\begin{aligned}
\sigma^1 &= \sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i(-1)^{p+1}\sum_{F\in\mathcal{F}_i(\Delta)}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q} \\
&= \sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i(-1)^{p+1}f_i(\Delta)\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}.
\end{aligned}$$

Furthermore, $\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q} = (-1)^{p+1}\binom{i}{p+1}$, and thus,

$$\sigma^1 = \sum_{p=0}^{d-1}\sum_{i=p+1}^{d}(-1)^i\binom{i}{p+1}f_i(\Delta).$$

□

For any $p \geq i$, $\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1} = \sum_{q=0}^{i+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1}$ which is zero by Lemma 6 since $l \leq i+1$. Thus one gets

$$\sigma_{l,i+1} = \sum_{p=0}^{i-1}(-1)^i(-1)^{p+1}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1}. \tag{2.2}$$

□

We can then prove the following lemma.

Lemma 4. *For $2 \leq l \leq i+1$, one has the following equalities*

$$\begin{aligned}
\sigma_{l,i+2} &= -2\sigma_{l,i+1}, \\
\chi_{l,i+2} &= -2\chi_{l,i+1}.
\end{aligned}$$

Proof. To prove the first equality, write $\binom{i+2}{q} = \binom{i+1}{q-1} + \binom{i+1}{q}$ to get

$$\begin{aligned}
\sigma_{l,i+2} = (-1)^{i+1}&\sum_{p=0}^{i}(-1)^{p+1}\sum_{q=1}^{p+1}(-1)^q\binom{i+1}{q-1}(p+1-q)^{l-1} \\
&+(-1)^{i+1}\sum_{p=0}^{i}(-1)^{p+1}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1}.
\end{aligned}$$

Notice that $\sum_{q=1}^{1}(-1)^q\binom{i+1}{q-1}(p+1-q)^{l-1}$ is 0. Similarly, in the second term, $\sum_{q=0}^{i+1}(-1)^q\binom{i+1}{q}(i+1-q)^{l-1}$ makes no contribution, by Lemma 6.

Then,

$$\begin{aligned}
\sigma_{l,i+2} = (-1)^{i+1}&\sum_{p=1}^{i}(-1)^{p+1}\sum_{q=1}^{p+1}(-1)^q\binom{i+1}{q-1}(p+1-q)^{l-1} \\
&+(-1)^{i+1}\sum_{p=0}^{i-1}(-1)^{p+1}\sum_{q=0}^{p+1}(-1)^q\binom{i+1}{q}(p+1-q)^{l-1}.
\end{aligned}$$

So, with the changes of indices $c = q - 1$ and $r = p - 1$ one gets

$$\begin{aligned}
\sigma_{l,i+2} &= (-1)^{i+1}\sum_{r=0}^{i-1}(-1)^r\sum_{c=0}^{r+1}(-1)^{c+1}\binom{i+1}{c}(r+1-c)^{l-1}\\
&\quad +(-1)^{i+1}\sum_{p=0}^{i-1}(-1)^{p+1}\sum_{q=0}^{p+1}(-1)^{q}\binom{i+1}{q}(p+1-q)^{l-1}\\
&= -2\sigma_{l,i+1}.
\end{aligned}$$

To prove the second equality stated in the lemma, use Lemma 6 to get rid of the term with $k = i + 1$ in the expression of $\chi_{l,i+2}$:

$$\begin{aligned}
\chi_{l,i+2} &= (-1)^{i-l+2}\sum_{k=1}^{i+1}\frac{(2^{i+1}-2^{i+1-k})}{k+1}\sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l\\
&= 2(-1)^{i-l}\sum_{k=1}^{i}\frac{(2^{i}-2^{i-k})}{k+1}\sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l\,,
\end{aligned}$$

which finishes the proof. □

Lemma 5. *For* $2 \le l \le i + 1 \le d + 1$, *we have* $\chi_{l,i+1} = \sigma_{l,i+1}$.

Proof. By Lemma 4 it is enough to prove the equality $\chi_{l,l} = \sigma_{l,l}$.

By Formula (2.2) one has,

$$\sigma_{l,l} = \sum_{p=0}^{l-2}(-1)^{l-1}\sum_{q=0}^{p+1}(-1)^{p+1-q}\binom{l}{q}(p+1-q)^{l-1} \tag{2.3}$$

$$= \sum_{b=0}^{l}(-1)^{l-1}\sum_{q=0}^{l-b}(-1)^{b}\binom{l}{q}(b)^{l-1} \tag{2.4}$$

$$= (-1)^{l-1}\sum_{b=1}^{l}\sum_{j=b}^{l}(-1)^{b}\binom{l}{j}b^{l-1}, \tag{2.5}$$

where we pass from (2.3) to (2.4) by setting $b = p + 1 - q$ and from (2.4) to (2.5) by putting $j = l - q$.

One sees that

$$\begin{aligned}
\chi_{l,l} &= \sum_{k=1}^{l-1}\frac{2^{l-1}-2^{l-1-k}}{k+1}\sum_{m=0}^{k+1}(-1)^m\binom{k+1}{m}m^l\\
&= \sum_{k=0}^{l-1}\frac{2^{l-1}-2^{l-1-k}}{k+1}\sum_{m=1}^{k+1}(-1)^m\binom{k+1}{m}m^l\,,
\end{aligned}$$

just noticing that the change of range does not affect the sum. Since $\binom{k+1}{m}\frac{m^l}{k+1} = \binom{k}{m-1}m^{l-1}$,

$$\begin{aligned}
\chi_{l,l} &= \sum_{k=0}^{l-1}(2^{l-1}-2^{l-1-k})\sum_{m=1}^{k+1}(-1)^m\binom{k}{m-1}m^{l-1}\\
&= 2^{l-1}\sum_{k=0}^{l-1}\sum_{m=1}^{k+1}(-1)^m\binom{k}{m-1}m^{l-1}-\sum_{k=0}^{l-1}2^{l-1-k}\sum_{m=1}^{k+1}(-1)^m\binom{k}{m-1}m^{l-1}\\
&= 2^{l-1}\sum_{m=1}^{l}(-1)^m m^{l-1}\sum_{k=m-1}^{l-1}\binom{k}{m-1}-\sum_{m=1}^{l}(-1)^m m^{l-1}\sum_{k=m-1}^{l-1}2^{l-1-k}\binom{k}{m-1}\\
&= 2^{l-1}\sum_{m=1}^{l}(-1)^m m^{l-1}\binom{l}{m}-\sum_{m=1}^{l}(-1)^m m^{l-1}\sum_{j=m}^{l}\binom{l}{j},
\end{aligned}$$

since $\sum_{k=m-1}^{l-1}\binom{k}{m-1}=\binom{l}{m}$ and $\sum_{k=m-1}^{l-1}2^{l-1-k}\binom{k}{m-1}$ by Lemma 7. By Lemma 6, $\sum_{m=1}^{l}(-1)^m m^{l-1}\binom{l}{m}=0$ thus

$$\chi_{l,l}=-\sum_{m=1}^{l}(-1)^m m^{l-1}\sum_{j=m}^{l}\binom{l}{j}$$

and $\chi_{l,l}=(-1)^l\sigma_{l,l}$. This is the desired equality for l even. For an odd $l>2$ we use the symmetry of the expression of $\sigma_{l,l}$ and write

$$\begin{aligned}
\sigma_{l,l}=\sum_{p=0}^{\frac{l-3}{2}}\Big(&(-1)^{p+1}\sum_{q=0}^{p+1}(-1)^q\binom{l}{q}(p+1-q)^{l-1}\\
&+(-1)^{l-1-p}\sum_{r=0}^{l-1-p}(-1)^r\binom{l}{r}(l-1-p-r)^{l-1}\Big).
\end{aligned}$$

With the change of index $q=l-r$ in the second term of the sum and noticing that the contribution for $q=p+1$ is zero, one gets

$$\sigma_{l,l}=\sum_{p=0}^{\frac{l-3}{2}}(-1)^{p+1}\sum_{q=0}^{l}(-1)^q\binom{l}{q}(p+1-q)^{l-1}.$$

The right hand side of the last equality is zero by Lemma 6, and thus, $\sigma_{l,l}=\chi_{l,l}=0$. □

According to Lemma 5 the coefficients of $\sigma(Z)$ and $\chi(\mathbb{R}Z)$ in the expressions of Lemma 2 are equal, and thus, $\sigma(Z)=\chi(\mathbb{R}Z)$. □

3. NAKAJIMA POLYTOPES AND M-SURFACES

In this section we give examples of families of M-surfaces obtained by T-construction as hypersurfaces of 3-dimensional toric varieties.

3.1. **Newton polytopes without maximal hypersurfaces.** We first explain that Itenberg-Viro's theorem (see [IV05]) on the existence of M-hypersurfaces of any degree in the projective spaces of any dimension cannot be generalized straightforwardly to all projective toric varieties. Indeed, in dimensions greater or equal to 3, there exist polytopes Δ such that no hypersurface in X_Δ with Newton polytope Δ is maximal (See Proposition 3.2). Note that this does not mean that the toric variety X_Δ does not admit M-hypersurfaces.

Clearly, if Δ is an interval $[a,b]$ in $\mathbb{R}$, where a and b are nonnegative integers, then there exists a maximal 0-dimensional subvariety in $\mathbb{C}P^1 = X_\Delta$ with the Newton polygon Δ.

If Δ is a polygon in the first quadrant of $\mathbb{R}^2$, then again there exists a maximal curve in X_Δ with the Newton polygon Δ. Such a curve can be constructed by the combinatorial patchworking: it suffices to take as initial data a primitive convex triangulation of Δ equipped with the following distribution of signs: an integer point (i,j) of Δ gets the sign "-" if i and j are both even, and gets the sign "+", otherwise (cf., for example, [Ite95], [IV96], [Haa98]).

However, in dimension 3 there are polytopes Δ such that no surface in X_Δ with the Newton polytope Δ is maximal (See Proposition 3.1).

Let k be a positive integer number, and Δ_k be the tetrahedron in $\mathbb{R}^3$ with vertices $(0,0,0), (1,0,0), (0,1,0)$, and $(1,1,k)$. Note that the only integer points of Δ_k are its vertices.



Figure 7. Tetrahedron Δ_3.

Proposition 3.1. *For any odd $k \geq 3$ and any even $k \geq 8$, there is no maximal surface in X_{Δ_k} with the Newton polytope Δ_k.*

The proof of the above Proposition can be found in [Ber03] and [Ber06]. There, this family of examples is generalized to higher dimensional polytopes and the following proposition is proved.

Proposition 3.2. *For any integer $d \geq 3$ there exist d-dimensional polytopes Δ such that no hypersurface in X_Δ with the Newton polytope Δ is maximal.*

3.2. **Construction of M-surfaces.** Using Corollary 2.2 of Theorem 2.1 we prove the existence of maximal surfaces in all nonsingular toric varieties corresponding to Nakajima polytopes of dimension 3 (see Figure 8).

A Nakajima polytope (See Introduction) is a ***nondegenerate*** if it is 0-dimensional or if $\bar{\Delta}$ is nondegenerate and f is positive on $\bar{\Delta}$.



FIGURE 8. Nakajima polytopes of dimension 2 and 3.



FIGURE 9. Slicing a nondegenerate Nakajima polytope.

Theorem 3.3. *Let Δ be a 3-dimensional Nakajima polytope corresponding to a nonsingular toric variety X_Δ. Then, there exists a maximal surface in X_Δ with the Newton polytope Δ.*

Proof. Let us first assume that Δ is nondegenerate.

Decompose Δ into slices $T_k = \Delta \cap \{(x, y, z) \in (\mathbb{R}^+)^3, k-1 \leq x \leq k\}$. Let s_k be the section $\Delta \cap \{(x, y, z) \in (\mathbb{R}^+)^3, x = k\}$. Denote by $B^{0,k}$, $B^{1,k}$, $B^{2,k}$, and $B^{3,k}$ the vertices of s_k (see Figure 9). We subdivide the slices T_k in two cones and two joins. Take the cones of apex $B^{0,2k}$ and $B^{0,2k+2}$ on s_{2k+1} and the cones of apex $B^{3,2k-1}$ and $B^{3,2k+1}$ on s_{2k}. Take primitive convex triangulations of the sections s_k. They induce a primitive triangulation of the cones. The joins $[B^{0,2k}, B^{1,2k}] * [B^{1,2k+1}, B^{3,2k+1}]$, $[B^{0,2k}, B^{2,2k}] * [B^{2,2k+1}, B^{3,2k+1}]$, $[B^{0,2k+2}, B^{1,2k+2}] * [B^{1,2k+1}, B^{3,2k+1}]$, $[B^{0,2k+2}, B^{2,2k+2}] * [B^{2,2k+1}, B^{3,2k+1}]$ are also naturally primitively triangulated which gives a primitive convex triangulation of Δ. Take the following distribution of signs at the integer points of Δ:

any integer point (i, j, k) gets "-" if j and k are both odd, and it gets "+", otherwise.

The polytope Δ is now equipped with a convex primitive triangulation and a sign distribution, so we can apply the T-construction as in Section 1.2.

Let p be an integer interior point of Δ and $\mathrm{Star}(p)$ be its star. Assume that p belongs to a section s_l. Then, $\mathrm{Star}(p)$ has two vertices c_1 and c_2 outside s_l. They are the apices of the two cones over s_l. They have the same parity (i.e. their coordinates have the same reduction modulo 2) and the distribution of signs depends only on the parity thus, in each octant, their symmetric copies carry the same sign. Consider an octant where the symmetric copy q_0 of p is isolated in s_l (i.e. all vertices of $\mathrm{Star}(q_0) \cap s_l$ except q_0 carry the sign opposite to the sign of q_0). Then, either in this octant the symmetric copies of c_1 and c_2 carry the sign opposite to the sign of q_0 (and, hence, q_0 is isolated), or $r(q_0)$ is isolated, where r is the reflection with respect to the coordinate plane $x = 0$. Thus, for each integer interior point p of Δ, there exists a symmetric copy q of p such that q is surrounded by a sphere $S^2(p) = \mathrm{Star}(q) \cap H^*$ (see Remark 1). Moreover at least one component of H^* intersects the coordinate planes. Thus, the T-surface constructed has at least $l^*(\Delta) + 1$ connected components, and Corollary 2.2 shows that this surface is maximal.



FIGURE 10. The star of p.

The degenerate case splits into two subcases (Since we only consider here Nakajima polytopes corresponding to nonsingular toric varieties). Either the Nakajima polytope is a truncated cylinder over a triangle corresponding to the projective plane, or it is a tetrahedron corresponding to the projective 3-space. The existence of M-surfaces in the latter case was proved by Viro in [Vir79]. In the former case the Nakajima polytope Δ is the convex hull of the triangles $((0, 0, 0), (m, 0, 0), (0, m, 0))$ and $((0, 0, l), (0, m, l + me), (m, 0, l + mf))$ for some integers m, l, e and f (see Figure 11).

Decompose Δ into slices $T_k = \Delta \cap \{(x, y, z) \in (\mathbb{R}^+)^3, k - 1 \leq x \leq k\}$. Let s_k be the section $\Delta \cap \{(x, y, z) \in (\mathbb{R}^+)^3, x = k\}$. We triangulate $\Delta \cap \{(x, y, z) \in (\mathbb{R}^+)^3, y \leq m - 1\}$ using the same triangulation as in the nondegenerate case. Take the cone over



Figure 11. A degenerate Nakajima polytope.

s_{m-1} of apex $(m, 0, 0)$ if m is even or $(m, 0, l + mf)$, otherwise. Take in the cone the triangulation induced by the triangulation of s_{m-1}. Subdivide the only remaining nonprimitive tetrahedron into primitive ones in the unique possible way. Take the same distribution of signs that in the nondegenerate case.

Then, as in the nondegenerate case, any interior point has an isolated symmetric copy. There is also a component intersecting the coordinate planes. Thus, the surface is maximal. □

Note that Theorem 3.3 produces, in particular, M-surfaces in $\Sigma_\alpha \times \mathbb{C}P^1$.

For a nonnegative integer α and positive integers l, m and n let $\delta_\alpha^{m,n}$ and $\Delta_\alpha^{l,m,n}$ be the polytopes defined in Section 1.4. Recall that the toric variety $X_{\Delta_\alpha^{l,m,n}}$ is isomorphic to $\Sigma_\alpha \times \mathbb{C}P^1$. Since $\Delta_\alpha^{l,m,n}$ is a Nakajima polytope, the following statement is a corollary of Theorem 3.3.

Corollary 3.4. *For any nonnegative integer α and any positive integers m, n and l, there exists a maximal surface in $\Sigma_\alpha \times \mathbb{C}P^1$ with the Newton polytope $\Delta_\alpha^{l,m,n}$.* □

4. Appendix

4.1. Usefull combinatorial lemmae.

We insert here the proofs of lemmae that we use quite often, mainly in Section 2.

Lemma 6. *Let l and i be nonnegative integers. Then, for $l+1 \leq i$, one has (See* [VLW92] *p. 71)*

$$\sum_{q=0}^{i} (-1)^q \binom{i}{q} q^l = \sum_{q=0}^{i} (-1)^q \binom{i}{q} (i-q)^l = 0 \tag{4.1}$$

and, as a consequence, for any integer p,

$$\sum_{q=0}^{i}(-1)^q\binom{i}{q}(p-q)^l = 0. \tag{4.2}$$

Proof. Formula 4.1 can be found in [VLW92] p. 71. We prove Equality (4.2) using (4.1).
Write $p-q=(i-q)+(p-i)$ and then,

$$(p-q)^l = \sum_{m=0}^{l}\binom{l}{m}(i-q)^{l-m}(p-i)^m.$$

So,

$$\begin{aligned}\sum_{q=0}^{i}(-1)^q\binom{i}{q}(p-q)^l &= \sum_{q=0}^{i}(-1)^q\binom{i}{q}\sum_{m=0}^{l}\binom{l}{m}(i-q)^{l-m}(p-i)^m \\ &= \sum_{m=0}^{l}\binom{l}{m}(p-i)^m\left(\sum_{q=0}^{i}(-1)^q\binom{i}{q}(i-q)^{l-m}\right)\end{aligned}$$

The last term is zero by Equality 4.1. □

Proof of lemma 1. From the definitions we see that

$$Q_\Delta(t) = (\sum_{k=0}^{d+1}\binom{d+1}{k}t^k(-1)^k)\cdot\sum_{n=0}^{\infty}(\sum_{i=0}^{d}a_i n^i)t^n,$$

and thus, $\Psi_j = \sum_{i=0}^{d}(\sum_{n=0}^{j}(-1)^{j-n}\binom{d+1}{j-n}n^i)a_i^\Delta$. Let $A_j = \sum_{n=0}^{j}(-1)^{j-n}\binom{d+1}{j-n}n^i$.
If $j \geq d+1$, then

$$A_j = \sum_{k=0}^{d+1}(-1)^k\binom{d+1}{k}(j-k)^i,$$

which is zero according to Lemma 6 Equality 4.2. □

Lemma 7. *One has* $\sum_{n=0}^{p}2^{p-n}\binom{n}{k} = \sum_{l=k+1}^{p+1}\binom{p+1}{l}$.

Proof. We use the fact that $\sum_{n=k}^{p}\binom{n}{k}=\binom{p+1}{k+1}$ and write

$$\begin{aligned}\sum_{n=0}^{p}2^{p-n}\binom{n}{k} &= \sum_{n=k}^{p}2^{p-n}\binom{n}{k}\\ &= \sum_{n=k}^{p}\binom{n}{k}+\sum_{i=0}^{p}(2^i)\sum_{n=k}^{p-i}\binom{n}{k}\\ &= \binom{p+1}{k+1}+\sum_{i=0}^{p-k}(2^i)\sum_{i=0}^{p-k}\binom{p+1-i}{k+1}\\ &= \binom{p+1}{k+1}+\sum_{n=k+1}^{p+1}2^{p+1-n}\binom{n}{k+1},\end{aligned}$$

and the result follows by induction. □

4.2. **Formula for the number of k-simplices.** In order to prove the formula of Proposition 1.7 we first check that it is true for all primitive simplices.

4.2.1. *Formula for a primitive simplex.* Let s_i be a primitive simplex of dimension i. Then

$$Ehr_{s_i}(\lambda)=\frac{1}{i!}(\lambda+1)\dots(\lambda+i)=\frac{1}{i!}\sum_{j=1}^{i+1}(-1)^{i+1-j}S_1(i+1,j)\lambda^{j-1},$$

where S_1 is the first Stirling number defined by the formula $\sum_{m=0}^{n}S_1(n,m)x^m=x(x-1)\dots(x-n+1)$ (see [BV97]). Thus

$$a_{j-1}^{s_i}=\frac{(-1)^{i+1-j}}{i!}S_1(i+1,j).$$

It remains to show that

$$N_r^{s_i}=\frac{r!}{i!}\sum_{l=r+1}^{i+1}S_2(l,r+1).S_1(i+1,l).$$

We have $\sum_{l=r+1}^{i+1}S_2(l,r+1).S_1(i+1,l)=\delta_{r,i}$ where $\delta_{r,i}$ is the Kronecker index (see [VLW92] p. 107), and thus,

$$\frac{r!}{i!}\sum_{l=r+1}^{i+1}S_2(l,r+1).S_1(i+1,l)=\delta_{r,i}$$

which is exactly the number of r-dimensional simplices contained in the interior of s_i (zero if $r\neq i$ and one othewise).

4.2.2. *Proof for a general polytope.*

Definition 4.1. *A triangulation τ of a polytope of dimension d is called* ***shellable*** *if there exists a numbering of its d-simplices $s_1, s_2, \dots, s_k$ such that for $i \in \{2, \dots, k\}$*

$$s_i \cap \cup_{j=1}^{i-1} s_j$$

is a nonempty union of $(d-1)$-simplices of τ homeomorphic to a $(d-1)$-ball. This numbering is called a shelling of τ.

Theorem 4.1 (Ziegler ([Zie95], p. 243))**.** *Every convex triangulation of a polytope is shellable.*

The formula of Proposition 1.7 holds for a point (with the convention that the point is in its interior). We now assume that the formula is true in all dimensions less than d. Let Δ be a d-dimensional polytope, and τ be a primitive triangulation of Δ. Fix $s_1, s_2, \dots, s_t$ a shelling of τ, and put $U_j = \cup_{i=1}^{j} s_j$. Note that U_j need not to be convex. Assume that the formula is true for U_j. Note that the formula also holds for $U_j \cap s_{j+1}$ which is $(d-1)$-dimensional, eventhough it is not a polytope. We know that the formula is also true for s_{j+1}.

Lemma 8. *The numbers N_r satisfy the relation*

$$N_r^{U_j} + N_r^{s_{j+1}} + N_r^{U_j \cap s_{j+1}} = N_r^{U_{j+1}}.$$

Proof. This follows immediately from the fact that $\overset{\circ}{U_j} \sqcup \overset{\circ}{s}_{j+1} \sqcup (U_j \overset{\circ}{\cap} s_{j+1}) = \overset{\circ}{U}_{j+1}$, where $\overset{\circ}{U}$ stands for the interior of U. $\square$

By induction hypothesis,

$$N_r^{U_j \cap s_{j+1}} \quad = \quad \sum_{l=r+1}^{d} \mathrm{N}(r+1, l)(-1)^{d-l} a_{l-1}^{U_j \cap s_{j+1}},$$

and since $a_d^{U_j \cap s_{j+1}} = 0$, we can also write

$$N_r^{U_j \cap s_{j+1}} \quad = \quad \sum_{l=r+1}^{d+1} \mathrm{N}(r+1, l)(-1)^{d-l} a_{l-1}^{U_j \cap s_{j+1}}.$$

Then $N_r^{U_{j+1}} = \sum_{l=r+1}^{d+1} \mathrm{N}(r+1, l)(-1)^{d-l+1}.(a_{l-1}^{U_j} + a_{l-1}^{s_{j+1}} - a_{l-1}^{U_j \cap s_{j+1}})$, and $a_{l-1}^{U_j} + a_{l-1}^{s_{j+1}} - a_{l-1}^{U_j \cap s_{j+1}}$ is precisely the coefficient $a_{l-1}^{U_{j+1}}$ in the Ehrhart polynomial of U_{j+1} (see [Ehr67] for Ehrhart polynomials of general polyhedra). This completes the proof of the formula. $\square$

REFERENCES

[Ber03] Benoît Bertrand. Ph.d. thesis: Maximal hypersurfaces and complete intersections in toric varieties. 2003. http://www.unige.ch/math/folks/bertrand.

[Ber06] Benoît Bertrand. Asymptotically maximal families of hypersurfaces in toric varieties. *Geom. Dedicata*, 118:49–70, 2006.

[Bri94] M. Brion. Points entiers dans les polytopes convexes. *Astérisque 227 , Séminaire Bourbaki 780*, 1994.

[BV97] Michel Brion and Michèle Vergne. Lattice points in simple polytopes. *J. Amer. Math. Soc.*, 10(2):371–392, 1997.

[Dai00] D. Dais. Über unimodulare, kohärente Triangulierungen von Gitterpolytopen. Beispiele und Anwendungen. *lecture note of the talk at the Grenoble summer school : Géométrie des variétés toriques 2000. 19 pages*, 2000.

[Dan78] V. I. Danilov. The geometry of toric varieties. *Uspekhi Mat. Nauk*, 33(2(200)):85–134, 247, 1978.

[DHZ01] Dimitrios I. Dais, Christian Haase, and Günter M. Ziegler. All toric local complete intersection singularities admit projective crepant resolutions. *Tohoku Math. J. (2)*, 53(1):95–107, 2001.

[DK87] V. Danilov and A. Khovanskii. Newton polyhedra and an algorithm for computing Hodge-Deligne numbers. *Math. USSR Izvestiya vol 29 No 2*, pages 279–298, 1987.

[Ehr67] E. Ehrhart. Sur un problème de géométrie diophantienne linéaire. I. Polyèdres et réseaux. *J. Reine Angew. Math.*, 226:1–29, 1967.

[Ehr94] E. Ehrhart. Un théorème arithmo-géométrique et ses généralisations. *L'ouvert n. 77*, pages 33–34, 1994.

[Ful93] W. Fulton. *Introduction to toric varieties.* Princeton University Press, 1993.

[GH78] P. Griffiths and J. Harris. *Principles of algebraic geometry.* John Willey and Sons, 1978.

[GKZ94] I. Gelfand, M. Kapranov, and A. Zelevinsky. *Discriminants, Resultants and Multidimensional Determinants.* Springer-Verlag, 1994.

[Haa98] Bertrand Haas. *Ph. D. thesis.* 1998.

[Ite95] Ilia Itenberg. Counter-examples to Ragsdale conjecture and T-curves. In *Real algebraic geometry and topology (East Lansing, MI, 1993)*, volume 182 of *Contemp. Math.*, pages 55–72. Amer. Math. Soc., Providence, RI, 1995.

[Ite97] I. Itenberg. Topology of real algebraic T-surfaces. *Revista mathematica univ. complutense de Madrid*, 10, no. supp.:131–152, 1997.

[IV96] Ilia Itenberg and Oleg Viro. Patchworking algebraic curves disproves the Ragsdale conjecture. *Math. Intelligencer*, 18(4):19–28, 1996.

[IV05] I. Itenberg and O. Viro. Maximal real algebraic hypersurface of projective space. *in preparation*, 2005.

[Mik00] G. Mikhalkin. Real algebraic curves, the moment map and amoebas. *Ann. of Math. (2)*, 151(1):309–326, 2000.

[Mik04] Grigory Mikhalkin. Amoebas of algebraic varieties and tropical geometry. In *Different faces of geometry*, Int. Math. Ser. (N. Y.), pages 257–300. Kluwer/Plenum, New York, 2004.

[Mik05] Grigory Mikhalkin. Enumerative tropical algebraic geometry in $\mathbb{R}^2$. *J. Amer. Math. Soc.*, 18(2):313–377 (electronic), 2005.

[Stu94] Bernd Sturmfels. Viro's theorem for complete intersections. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 21(3):377–386, 1994.

[Vir79] O. Ja. Viro. Construction of M-surfaces. *Funktsional. Anal. i Prilozhen.*, 13(3):71–72, 1979.

[Vir84] Oleg Viro. Gluing of plane algebraic curves and construction of curves of degree 6 and 7 (lnm 1060). *Lecture Notes in Mathematics*, pages 187–200, 1984.

[Vir01] Oleg Viro. Dequantization of real algebraic geometry on logarithmic paper. In *European Congress of Mathematics, Vol. I (Barcelona, 2000)*, volume 201 of *Progr. Math.*, pages 135–146. Birkhäuser, Basel, 2001.

[Vir04] O. Viro. Patchworking real algebraic varieties. *preprint Uppsala University*, 2004.

[VLW92] J. H. Van Lint and R. M. Wilson. *A Course in Combinatorics*. Cambridge university press, 1992.

[Zie95] G. Ziegler. *Lectures on polytopes*. Springer-Verlag, 1995.

Section de mathématiques, Université de Genève, case postale 64, 2-4 rue du lièvre, 1211 Genève 4, Suisse

E-mail address: `benoit.bertrand@math.unige.ch`

URL: `http://name.math.univ-rennes1.fr/benoit.bertrand`